

Combaining ability and heterosis in sorghum_Sabrao

by Sulistyawati Sulistyawati

Submission date: 20-Mar-2023 01:26AM (UTC-0400)

Submission ID: 2041409892

File name: 2022-2_Combaining_ability_and_heterosis_in_sorghum_Sabrao.pdf (584.35K)

Word count: 8353

Character count: 44770

1
 SABRAO Journal of Breeding and Genetics
 54 (1) 30-43, 2022
<http://doi.org/10.54910/sabrao2022.54.1.4>
<http://sabraojournal.org/>
 pISSN 1029-7073; eISSN 2224-8978



39
COMBINING ABILITY AND HETEROSIS IN SORGHUM (*Sorghum bicolor* L.)

**MAFTUCHAH^{1,2*}, H. WIDYANINGRUM¹, A. ZAINUDIN^{1,3}, SULISTYAWATI³,
 H.A. RESWARI², and H. SULISTIYANTO²**

6
¹Department of Agrotechnology, University of Muhammadiyah Malang, Malang, East Java, Indonesia

²Center for Biotechnology, University of Muhammadiyah Malang, Malang, East Java, Indonesia

³Faculty of Agriculture, Merdeka 'Pasuruan' University, 'Pasuruan' City, East Java, Indonesia

*Corresponding author email: maftuchah@umm.ac.id

Email addresses of co-authors: halimahwidya9797@gmail.com, aguszainudinumm@gmail.com,
 mommyandri@gmail.com, helvi_ardana@umm.ac.id, hendrysulistiyanto21@gmail.com

1
SUMMARY

For the sustainable improvement of sorghum (*Sorghum bicolor* L.), local genotypes were conventionally crossed in East Java, Indonesia to determine a) the crossing success of different genotypes, b) germination percentage and seed setting traits, and c) the combining ability and heterotic effect of sorghum parental genotypes and their F₁ diallel hybrids. Three sorghum genotypes ('Lamongan-1', 'Tulungagung-2', and 'Jombang') from the local germplasm were collected from various regions in East Java, Indonesia, and were crossed in a complete diallel design. The experiment was carried out in a randomized complete block design with four replications during 2018–2019 at the Agrotechnology Laboratory, Faculty of Agriculture, University of Muhammadiyah, Malang, Indonesia. Results showed that the parental genotypes and their F₁ hybrids exhibited significant differences in seed weight and seed diameter and nonsignificant differences in germination percentage and seed setting. The parental genotype 'Tulungagung-2' and its hybrids 'Jombang' × 'Tulungagung-2', 'Tulungagung-2' × 'Lamongan-1', and 'Jombang' × 'Tulungagung-2' presented the highest germination percentages, seed setting, seed weights, and seed diameters. The genotype 'Tulungagung-2' had the highest general combining ability and was identified as a good general paternal and maternal parent for the production of promising F₁ hybrids. The hybrid 'Jombang' × 'Lamongan-1' had the highest specific combining ability for all of the characters and was recognized as a promising F₁ hybrid for improving sorghum yield. Cluster analysis divided all of the parental genotypes and their F₁ hybrids into two large groups with a similarity of 13.13%.

Keywords: Conventional breeding, parental genotypes, F₁ diallel hybrids, crossing success, germination percentage, heterosis, heterobeltiosis, combining ability

Key findings: 'Tulungagung-2' was identified as a good general paternal and maternal parent for the production of promising F₁ hybrids with the highest germination percentage, seed setting, seed weight, and seed diameter. The F₁ hybrid 'Jombang' × 'Lamongan-1' was identified as the best specific combiner. The genotype 'Tulungagung-2' can be recommended for the assembly of superior sorghum cultivars with high productivity.

1
 Communicating Editor: Dr. Kamile Ulukapi

Manuscript received: March 25, 2021; Accepted: February 16, 2022.

© Society for the Advancement of Breeding Research in Asia and Oceania (SABRAO) 2022

3
To cite this manuscript: Maftuchah, Widyaningrum H, Zainudin A, Sulistyawati, Reswari HA, Sulistiyanto H (2022). Combining ability and heterosis in Sorghum (*Sorghum bicolor* L.). *SABRAO J. Breed. Genet.* 54(1): 30-43. <http://doi.org/10.54910/sabrao2022.54.1.4>

INTRODUCTION

Sorghum (*Sorghum bicolor* L.) belongs to the family Gramineae; it is widely adapted and can be grown on marginal lands, such as dry, alkaline, acidic, or even infertile lands (Sembiring and Subekti, 2010). Sorghum is not only limited to use as food and feed ingredients, it can also be used as a processed material; moreover, given that its seeds contain ash (1.6%), fat (3.1%), protein (10.4%), carbohydrates (70.7%), crude fiber (2.0%), and energy (329 kcal), it can serve as a multipurpose crop (Suarni, 2012; Reddy *et al.*, 2014; Jimoh and Abdullahi, 2017). It is a versatile crop that is used as a food, feed, and industrial ingredient. As a food ingredient, sorghum is not considerably different from other cereals (ICRISAT, 2004). In general, the protein content of sorghum is higher than that of corn, brown rice, and barley but lower than that of wheat. The fat content of sorghum is higher than that of brown rice, wheat, and millet but lower than that of corn.

Sorghum is suitable for cultivation in Indonesia because this country has a tropical climate. It is relatively drought-tolerant and could be grown in areas with low soil fertility (Panjaitan *et al.*, 2015). In Indonesia, the past research on the development of sorghum is limited compared with that on other cereal crops, such as rice, maize, and wheat (Boonlertnirun *et al.*, 2012). In Indonesia, the level of production and quality of sorghum remain low, and many regions still rely on the cultivation of local genotypes.

In Indonesia, the lack of superior cultivars is the main problem in sorghum development because little work has been conducted on their identification, development, and characterization (Susilowati and Saliem, 2013; Rao *et al.*, 2016). Therefore, collecting and developing the local germplasm through different breeding programs is necessary to obtain wide genetic diversity in sorghum. The sorghum breeding program in East Java, Indonesia, is under development, and several local genotypes have been crossed to obtain superior genotypes. On the basis of the initial characterization of the local germplasm in East Java, eight sorghum genotypes ('Pasuruan', 'Lamongan-1', 'Lamongan-2', 'Tuban', 'Sampang-2', 'Tulungagung-1', 'Tulungagung-2', and 'Jombang') have been developed with increased potential and desirable agronomic characters and can be used as parental genotypes for the development of superior cultivars in future breeding programs (Sulistiyawati *et al.*, 2019a).

The sorghum genotype 'Lamongan-1' meets the specific criterion of maturation within 77 days (Sulistiyawati *et al.*, 2019b). The selection criteria for the development of superior sorghum cultivars include early maturity, medium plant height, and desirable protein and tannin contents (Subagio and Aqil, 2013). The sorghum genotype 'Lamongan-1' also has the highest leaf length and width, the highest number of seeds per panicle, and the second heaviest seed weight per panicle after the genotype 'Tulungagung-2'. The genotype 'Tulungagung' has the second highest plant height after the genotype 'Jombang', the highest fresh and dry 1000-seed weight with the second-highest number of seeds per panicle after the genotype 'Lamongan-1', but earlier flowering and harvesting than 'Lamongan-1' and 'Jombang'. Maftuchah *et al.* (2020, 2021) revealed that genotype 'Tulungagung-2' has more chances for development than other genotypes because it has the highest fresh and dry seed weights. The sorghum genotype 'Jombang' has the advantage of the highest plant height, leaf number, and reducing sugar content (Sulistiyawati *et al.*, 2019a, b).

The crossing success rate and seed germination of F₁ hybrids must to be determined to validate whether the F₁ derivatives can produce vigorous seeds such that selection can be made on the basis of the availability of the seeds of different cross combinations. Genetic parameters, such as heterosis and heterobeltiosis, are used to assess F₁ hybrids quantitatively (Boonlertnirun *et al.*, 2012). Heterosis and heterobeltiosis can be used to improve plant characters by identifying parental combinations that can produce high-yielding F₁ strains. In sorghum, the selection of promising parental genotypes to produce a superior hybrid is dependent on heterotic effects F₁ populations (Ringo *et al.*, 2015; Jaikishan *et al.*, 2019).

Poehlman (1983) reported that not all pure-line combinations will produce superior hybrids. Therefore, pure lines must be tested for their cohesiveness to determine the best combination for hybrid seed production. Welsh (1981) stated that the population with the highest general combining ability (GCA) effects has high chances to produce F₁ hybrids with the highest specific combining ability (SCA) values. The first step in determining the yield between the lines is to evaluate GCA and SCA. GCA is a general concept for classifying relatively pure strains in accordance with their hybrid's appearance (Hallauer and Miranda, 1988; Rini *et al.*, 2017). This information is

needed to obtain the results of parental combinations that will produce offspring with high potential from which high yields can be achieved if their offspring from their cross combinations have positive heterosis and high combining ability. Given the above considerations, the present study was planned to determine a) the crossing success of different sorghum genotypes, b) germination percentage and seed setting traits, and c) the combining ability and heterotic effects of sorghum parental genotypes and their F₁ diallel hybrids at East Java, Indonesia.

MATERIALS AND METHODS

Breeding material

The present research on sorghum was carried out during 2018–2019 at the Agrotechnology Laboratory, Faculty of Agriculture, University of Muhammadiyah Malang, Indonesia. Three sorghum genotypes ('Lamongan-1', 'Tulungagung-2', and 'Jombang') belonging to the local germplasm were collected from various regions in East Java, Indonesia. All of the sorghum genotypes were crossed in a complete diallel fashion at Purutrejo Village, Purworejo Subdistrict, Pasuruan City, Indonesia (–7°39'53.39" latitude, 112°54'17.38" east longitude).

Crossing program

Sorghum seeds were sown in polybags with dimensions of 4 cm × 3 cm and containing a mixture of soil + compost + rice husk. After emergence and 20 days of planting, the sorghum seedlings were transferred to the field, where each hill contained a single plant.

Emasculatation

The initial step was to prepare the equipment to be used in the crossing program of sorghum. Then, the sorghum flowers to be emasculated were selected. Flower selection was carried out as follows: two repetitions were used in one panicle, wherein for one repetition 3–4 secondary stalks were taken. Emasculatation was started from the base to the tip of the secondary stalk and carried out by opening the husks (glume) of the sorghum flowers on one side by using tweezers. Then, the tweezers were inserted into the husks to remove slowly the three stamens (anthers) in the sorghum flowers. Finally, the emasculated sorghum flowers were covered with paper

envelopes. The sorghum flowers were covered per repeat of the secondary flower stalk that has been emasculated. A large number of flowers were crossed. Specifically, 90 flowers for 'Lamongan-1' × 'Tulungagung-2', 91 flowers for 'Lamongan-1' × 'Jombang', 91 flowers for 'Tulungagung-2' × 'Lamongan-1', 84 flowers for 'Tulungagung-2' × 'Jombang', 81 flowers for 'Jombang' × 'Lamongan-1', and 84 flowers for 'Jombang' × 'Tulungagung-2' were crossed.

Pollination

The first step was to prepare the equipment for pollination. Then, the pollen grains of the selected genotypes were taken from stamens for use as male parents ('Lamongan-1', 'Tulungagung-2', and 'Jombang') by shaking the sorghum flower panicles slowly. Furthermore, pollination was carried out by opening the husks (glume) of sorghum flowers on one side of the husk by using tweezers and holding one side of the husk with fingers and then inserting the stamens vertically by using the tweezers into the husk and removing the other side of the chaff that was held with a finger to close the sorghum flower again. Then, the sorghum flowers were covered with paper bags. The secondary stalk of the pollinated sorghum flowers was also covered. The final stage was labeling information, i.e., the names of the parental genotypes used in the cross (female × male), replication, and the date of the cross.

Harvesting of the crosses seeds

The different crosses were harvested in accordance with the maturity and age of the sorghum genotypes. Harvest time was decided by looking at the visual characteristics of the stems, leaves, and panicles of ready-to-harvest sorghum. These visual characteristics included the yellowing of the leaves and stems, the formation of perfect panicles, and the hardening of the seeds. After harvesting, the sorghum was dried in the sun.

Sowing of parental genotypes and F₁ hybrid seeds

The different types of equipment and material used in the germination test were first prepared. The seeds of the parental genotypes and their crosses were soaked in warm water (40 °C) for 2 h, after which they were soaked in a fungicide solution for 15 min and then dried. Furthermore, the germination test was

carried out by using the interpaper test method by soaking three sheets of jaw paper and placing them in a plastic box, after which the tissue was placed on top of the jaw paper. The treated seeds of the sorghum genotypes were arranged on tissue paper, then covered with three sheets of moistened bamboo paper. The plastic boxes were closed and labeled with the date and genotype information and then placed at room temperature (28 °C).

Planting seeds

After the sprouting of the sorghum seeds and the appearance of the radicle, the candidate genotypes were immediately planted in 4 cm × 3 cm polybags containing a mixture of soil + compost + husk. At 20 days after sowing (DAS), the plants of the different sorghum genotypes were transferred to the planting area in the field, where each hole contained one sorghum seed at the depth of 5 cm. The said experiment was carried out in a randomized complete block design with six treatments and four replications.

Data recorded

The data on the different quantitative characters of sorghum parental genotypes and F₁ hybrids were recorded. These data included germination percentage (normal, abnormal, and dead seeds), seed setting, seed weight, and seed diameter. The germination percentage of the normal sprouts was observed and recorded from days 1 to 14. Kamil (1979) and Juliantisa (2017) stated that the criteria for normal sprouts are the presence of primary and secondary roots, long and short hypocotyls, and one primary leaf. Abnormal sprouts are those that do not meet the requirements for normal sprouts (DG-TPH, 2000). Data on abnormal sprouts were recorded from observations on the germination of a seed in which one or both essential elements (plumule and radicle) did not grow properly until the last day of observation. Dead seeds were those that at the end of the test were no longer hard or fresh and were usually moldy, completely rotted, and lacking the main elements for emergence. Data on dead seeds were recorded by counting the number of moldy and rotten seeds that did not germinate until the last day of observation.

Sprouts (%) = Number of live sprouts/number of seedlings × 100%

Germination power (%) = Number of sprouts/number of seeds grown × 100

In accordance with Juliantisa (2017), the sorghum germination percentage was calculated by using the following formula:

$$DB (\%) = \frac{\text{Normal sprouts}}{\text{Number of seeds}} \times 100$$

The seeds (%) were those seeds that formed after sorghum was harvested and dried. The formula for seed setting was as follows (Xie *et al.*, 2016):

$$\text{Seed set} = \frac{\text{Seeds formed}}{\text{Flowers crossed}} \times 100$$

Average seeds formed (#): Number of seeds formed in each cross/number of crosses

The average sorghum seed weight was recorded as follows:

$$\text{Average seed weight} = \frac{\text{Total seed weight}}{\text{number of seeds}}$$

Statistical analysis

All the obtained data were analyzed by using analysis of variance (ANOVA) to determine whether significant differences existed among the treatments or not (Steel *et al.*, 1997). Differences between treatments were revealed through the honestly significant difference (BNJ) test at the 5% level. Germination data were tested by analysis of means at the 5% level.

Heterosis and heterobeltiosis

After obtaining the data on the mean performance of the parental genotypes and their F₁ hybrids for various traits in sorghum, the heterotic effects over mid- and high-parents were calculated in accordance with Fehr (1987) as follows:

$$hMP (\%) = [(\mu F_1 - \mu MP) / \mu MP] \times 100$$

$$hHP (\%) = [(\mu F_1 - \mu HP) / \mu HP] \times 100$$

where

hMP = Average heterosis of parents (heterosis)

μMP = Mean value of both parents

hHP = Average heterosis of the highest parent (heterobeltiosis)

μ_{HP} = Mean value of the highest parent (high parent)

μ_{Fmean} = Value of the results of the study

Combining ability analysis

After significant differences were found among the parental inbred lines and their F_1 hybrids for various traits, the data were further subjected to combining ability analysis in accordance with Griffing (1956) Method-I based on Eisenhart's Model-II (Singh and Chaudhary, 1985) as follows

General combining ability

$$g_i = \frac{1}{2n} (Y_{i.} + Y_{.i}) - \frac{1}{n^2} Y_{..}$$

where

g_i = GCA effects for line i

n = Number of parents/varieties

$Y_{i.}$ = Total of mean values of F_1 s resulting from crossing i th lines with j th lines

$Y_{.i}$ = Total of mean values of F_1 s resulting from crossing i th lines with j th lines

$Y_{..}$ = Grand total of all the mean values in the table

Specific combining ability

$$s_{ij} = \frac{1}{2} (Y_{ij} + Y_{ji}) - \frac{1}{2n} (Y_{i.} + Y_{.i} + Y_{.j} + Y_{.j}) + \frac{1}{n^2} Y_{..}$$

where

s_{ij} = SCA between the i th and j th lines

Y_{ij} = Mean value of the F_1 resulting from crossing the i th and j th lines

\bar{y}_{ij} = Mean value for F_1 resulting from crossing the j th and i th varieties

$Y_{i.}$ = Total of mean values of F_1 's resulting from crossing j th line with i th varieties

\bar{y}_{ij} = Reciprocal values of Y_{ij}

$Y_{.j}$ = Total values for F_1 s resulting from crossing the i th line with j th line

$Y_{.j}$ = Values of reciprocal F_1 s of $Y_{.j}$

$Y_{..}$ = Grand total of the observations

Reciprocal effects

$$r_{ij} = \frac{1}{2} (Y_{ij} - Y_{ji})$$

where

r_{ij} = Reciprocal effects of i th and j th varieties/lines

n

Y_{ij} = Mean values for the F_1 resulting from crossing the i th and j th lines

Y_{ji} = Reciprocal values of F_1 resulting from Y_{ij}

Cluster analysis

Cluster analysis was also performed to determine the kinship relationship through the level of similarity between the crosses of three sorghum parental genotypes. The quantitative and qualitative data obtained from these sorghum parental genotypes and their crosses were analyzed.

RESULTS

Germination percentage

The germination percentages of the sorghum parental genotypes and their F_1 hybrids ranged from 76.00% to 92.00% and from 0.00% to 100%, respectively (Table 1). The parental genotypes showed higher germination percentages (86.67%) than their F_1 hybrids (68.85%). Among the parental genotypes, 'Tulungagung-2' and 'Jombang' showed the highest and the same germination (92.00%) rates. The lowest germination rate was shown by 'Lamongan-1' (76.00%). Among the F_1 hybrids, 'Lamongan-1' × 'Jombang', 'Tulungagung-2' × 'Lamongan-1', and 'Jombang' × 'Tulungagung-2' showed the highest germination percentages of 100%, 88.09%, and 83.33%, respectively. However, the F_1 hybrid 'Lamongan-1' × 'Tulungagung-2' exhibited zero germination percentage. The F_1 hybrids 'Jombang' × 'Lamongan-1' (75.00%) and 'Tulungagung-2' × 'Jombang' presented moderate germination percentage (66.67%).

Seed setting

Seed setting is an important and final output of every crop plant. The average seed setting of the sorghum F_1 hybrids ranged from 0.75 to 9.00 with percentages of 3.57% and 37.50%, respectively (Table 2). On average, the F_1 hybrid 'Tulungagung-2' × 'Lamongan-1' showed the maximum number of seeds (9.00) with a percentage of 37.50%, followed by the F_1 hybrid 'Jombang' × 'Tulungagung-2' (4.25 and 18.97%). The four F_1 hybrids 'Lamongan-1' × 'Jombang', 'Lamongan-1' × 'Tulungagung-2', 'Tulungagung-2' × 'Jombang', and 'Jombang' × 'Lamongan-1' presented the lowest number and percentage of seed setting with ranges of 0.75–1.50 and 3.57%–7.26%, respectively.

Table 1. Mean performance of sorghum parental genotypes and their F₁ hybrids for germination percentage.

| Genotypes | Average | | | | | Germination (%) |
|------------------------------|--------------|--------------|----------------|------------------|-----------------|-----------------|
| | Seedling (#) | Live sprouts | Normal sprouts | Abnormal sprouts | Succumb sprouts | |
| Parental genotypes | | | | | | |
| Lamongan-1 | 25 | 21 | 19 | 2 | 4.00 | 76.00 |
| Tulungagung-2 | 25 | 24 | 23 | 1 | 1.00 | 92.00 |
| Jombang | 25 | 24 | 23 | 1 | 1.00 | 92.00 |
| Mean | - | - | - | - | - | 86.67 |
| F₁ hybrids | | | | | | |
| Lamongan-1 × Tulungagung-2 | 2 | 0 | 0 | 0 | 2.00 | 0.00 |
| Lamongan-1 × Jombang | 2 | 2 | 2 | 0 | 0.00 | 100 |
| Tulungagung-2 × Lamongan-1 | 12 | 11 | 10 | 1 | 0.67 | 88.09 |
| Tulungagung-2 × Jombang | 2 | 2 | 1 | 0 | 0.50 | 66.67 |
| Jombang × Lamongan-1 | 3 | 3 | 3 | 0 | 0.50 | 75.00 |
| Jombang × Tulungagung-2 | 9 | 9 | 8 | 1 | 0.00 | 83.33 |
| Mean | - | - | - | - | - | 68.85 |

Table 2. Mean performance of sorghum F₁ diallel hybrids for seed setting.

| F ₁ hybrids | Replications | Flowers crossed (#) | Seeds formed (#) | Seeds formed (%) | Average seeds formed (#) | Average seeds formed (%) |
|----------------------------|--------------|---------------------|------------------|------------------|--------------------------|--------------------------|
| Lamongan-1 × Tulungagung-2 | 1 | 20 | 0 | 0.00 | 1.00 | 4.38 |
| | 2 | 20 | 1 | 5.00 | | |
| | 3 | 24 | 3 | 12.50 | | |
| | 4 | 26 | 0 | 0.00 | | |
| Lamongan-1 × Jombang | 1 | 25 | 0 | 0.00 | 0.75 | 3.57 |
| | 2 | 21 | 2 | 9.52 | | |
| | 3 | 21 | 1 | 4.76 | | |
| | 4 | 24 | 0 | 0.00 | | |
| Tulungagung-2 × Lamongan-1 | 1 | 25 | 9 | 36.00 | 9.00 | 37.50 |
| | 2 | 21 | 0 | 0.00 | | |
| | 3 | 25 | 21 | 84.00 | | |
| | 4 | 20 | 6 | 30.00 | | |
| Tulungagung-2 × Jombang | 1 | 22 | 1 | 4.55 | 1.00 | 4.89 |
| | 2 | 22 | 0 | 0.00 | | |
| | 3 | 20 | 0 | 0.00 | | |
| | 4 | 20 | 3 | 15.00 | | |
| Jombang × Lamongan-1 | 1 | 21 | 4 | 19.05 | 1.50 | 7.26 |
| | 2 | 20 | 0 | 0.00 | | |
| | 3 | 20 | 2 | 10.00 | | |
| | 4 | 20 | 0 | 0.00 | | |
| Jombang × Tulungagung-2 | 1 | 23 | 14 | 60.87 | 4.25 | 18.97 |
| | 2 | 21 | 0 | 0.00 | | |
| | 3 | 20 | 3 | 15.00 | | |
| | 4 | 20 | 0 | 0.00 | | |

Seed weight

The seed weight of the sorghum parental genotypes ranged from 0.023 g to 0.028 g and that of F₁ hybrids ranged from 0.008 g to 0.028 g (Table 3). Overall, the parental genotypes exhibited bolder seeds and higher

seed weight (0.025 g) than their F₁ hybrids (0.019 g). Among the parental genotypes, 'Tulungagung-2' (0.028 g) presented the maximum seed weight, followed by 'Lamongan-1' (0.025 g) and 'Jombang' (0.023 g). Among the F₁ hybrids, the highest seed weight was shown by the two F₁ hybrids

26

Table 3. Mean performance of sorghum parental genotypes and their F₁ hybrids for seed weight.

| Genotypes | Seed weight (g) |
|------------------------------|-----------------|
| Parental genotypes | |
| Lamongan-1 | 0.025 |
| Tulungagung-2 | 0.028 |
| Jombang | 0.023 |
| Mean | 0.025 |
| F₁ Hybrids | |
| Lamongan-1 × Tulungagung-2 | 0.008 |
| Lamongan-1 × Jombang | 0.019 |
| Tulungagung-2 × Lamongan-1 | 0.017 |
| Tulungagung-2 × Jombang | 0.019 |
| Jombang × Lamongan-1 | 0.020 |
| Jombang × Tulungagung-2 | 0.028 |
| Mean | 0.019 |

'Jombang' × 'Tulungagung-2' (0.028 g) and 'Jombang' × 'Lamongan-1' (0.020 g). However, the F₁ hybrid 'Lamongan-1' × 'Tulungagung-2' (0.008 g) showed shrunken seeds and the lowest seed weight among all F₁ hybrids and parental genotypes. All other F₁ hybrids showed moderate values for seed weight.

Seed diameter

The seed vertical and horizontal dimensions of the sorghum parental genotypes ranged from 0.20 cm to 0.28 cm and 0.15 cm to 0.18 cm, respectively (Table 4). The seed vertical and horizontal dimensions of the F₁ hybrids varied from 0.11 cm to 0.27 cm and from 0.08 cm to 0.18 cm, respectively. On average, the parental genotypes had larger seeds in terms of vertical and horizontal dimensions (0.26 and 0.16 cm) than F₁ hybrids (0.18 and 0.12 cm). Among parental genotypes, 'Tulungagung-2' showed the maximum values for seed vertical and horizontal dimensions (0.31 and 0.18 cm), followed by 'Lamongan-1' (0.28 and 0.15 cm), whereas the lowest seed vertical and horizontal dimensions were presented by 'Jombang' (0.20 and 0.15 cm). Among F₁ hybrids, 'Tulungagung-2' × 'Lamongan-1' exhibited the highest values for seed vertical and horizontal dimensions (0.27 and 0.18 cm) and 'Jombang' × 'Tulungagung-2' (0.25 and 0.15 cm). The F₁ hybrids 'Lamongan-1' × 'Tulungagung-2' (0.12 and 0.08 cm) and 'Lamongan-1' × 'Jombang' (0.11 and 0.08 cm) had close and the lowest seed vertical and horizontal dimensions. All other F₁ hybrids presented moderate values for seed vertical and horizontal dimensions. In sorghum, seed diameter is thought to be more influenced by genetic factors than by other factors.

Heterosis and heterobeltiosis

For germination percentage, the sorghum F₁ hybrid 'Lamongan-1' × 'Jombang' showed the highest positive heterotic effects over the mid- (0.19%) and better-parents (0.09%), followed by the F₁ hybrid 'Tulungagung-2' × 'Lamongan-1' (Table 5). However, all other F₁ hybrids exhibited negative heterosis and heterobeltiosis for germination percentage. The F₁ hybrid 'Jombang' × 'Tulungagung-2' exhibited the highest positive heterosis (0.10%) and heterobeltiosis (0.00%) for seed weight. However, all other F₁ hybrids revealed negative heterotic effects over their mid- and better-parents for seed weight. For seed vertical and horizontal dimensions, almost all of the F₁ hybrids, except for the F₁ hybrid 'Tulungagung-2' × 'Lamongan-1' with positive heterotic effects over its mid-parent (0.09%) and zero heterotic effect over its better parent for the said trait, exhibited negative heterotic effects over its mid- and high-parents.

Combining ability analysis

ANOVA showed that crosses ($P \leq 0.05$) and SCA ($P \leq 0.01$) mean squares were significant for seed vertical diameter (Table 6). The SCA mean squares were significant ($P \leq 0.05$) for vertical and horizontal diameter. However, all other mean squares for crosses, GCA, and SCA were nonsignificant for other traits. The sorghum genotype 'Tulungagung-2' had the highest positive GCA effects for seed weight and seed vertical and horizontal diameters, i.e., 0.0001, 0.0262, and 0.0164, respectively (Table 7). 'Lamongan-1' and 'Jombang' had the highest positive GCA effects for vertical diameter and seed weight, i.e., 0.0055 and 0.0002.

However, all the parental genotypes presented negative GCA effects for other traits. In the case of SCA effects in F₁ hybrids, the parental genotypes with the highest positive GCA

effects were also unable to produce hybrids with positive SCA effects for the said traits, and all the F₁ hybrids revealed negative SCA effects for the studied traits (Table 8).

26
Table 4. Mean performance of sorghum parental genotypes and their F₁ hybrids for seed diameter.

| Genotypes | Seed diameter (cm) | |
|----------------------------|--------------------|------------|
| | Vertical | Horizontal |
| Parental genotypes | | |
| Lamongan-1 | 0.28 | 0.15 |
| Tulungagung-2 | 0.31 | 0.18 |
| Jombang | 0.20 | 0.15 |
| Means | 0.26 | 0.16 |
| F ₁ Hybrids | | |
| Lamongan-1 × Tulungagung-2 | 0.12 | 0.08 |
| Lamongan-1 × Jombang | 0.11 | 0.08 |
| Tulungagung-2 × Lamongan-1 | 0.27 | 0.18 |
| Tulungagung-2 × Jombang | 0.14 | 0.10 |
| Jombang × Lamongan-1 | 0.16 | 0.11 |
| Jombang × Tulungagung-2 | 0.25 | 0.15 |
| Means | 0.18 | 0.12 |

4
Table 5. Sorghum F₁ hybrids with heterotic effects over their mid- and better-parents for germination percentage, seed weight, and seed diameter (vertical and horizontal).

| F ₁ Populations | Germination | | Seed weight | | Seed diameter (%) | | | |
|----------------------------|-------------|--------|-------------|--------|-------------------|--------|------------|--------|
| | 27 | | | | Vertical | | Horizontal | |
| | MP (%) | HP (%) | MP (%) | HP (%) | MP (%) | HP (%) | MP (%) | HP (%) |
| Lamongan-1 × Tulungagung-2 | -1.00 | -1.00 | -0.70 | -0.71 | -0.59 | -0.61 | -0.52 | -0.56 |
| Lamongan-1 × Jombang | 0.19 | 0.09 | -0.21 | -0.24 | -0.54 | -0.61 | -0.47 | -0.47 |
| Tulungagung-2 × Lamongan-1 | 0.05 | -0.04 | -0.36 | -0.39 | -0.08 | -0.13 | 0.09 | 0.00 |
| Tulungagung-2 × Jombang | -0.28 | -0.28 | -0.25 | -0.32 | -0.45 | -0.55 | -0.39 | -0.44 |
| Jombang × Lamongan-1 | -0.11 | -0.18 | -0.17 | -0.20 | -0.33 | -0.43 | -0.27 | -0.27 |
| Jombang × Tulungagung-2 | -0.09 | -0.09 | 0.10 | 0.00 | -0.02 | -0.19 | -0.09 | -0.17 |

MP: Mid-parent heterosis, HP: High-parent heterosis

4
Table 6. Analysis of the variance for a 3 × 3 sorghum diallel cross.

| Source of variation | d.f. | Mean squares | | |
|---------------------|------|----------------------|----------------------|----------------------|
| | | Weight per seed | DVB | DHB |
| Replications | 3 | 0.0001 ^{NS} | 0.0147 ^{NS} | 0.0047 ^{NS} |
| Crosses | 8 | 0.0002 ^{NS} | 0.0322* | 0.0118 ^{NS} |
| GCA | 2 | 0.0000 ^{NS} | 0.0206 ^{NS} | 0.0055 ^{NS} |
| SCA | 3 | 0.0004 ^{NS} | 0.0561** | 0.0207* |
| Reciprocals | 3 | 0.0001 ^{NS} | 0.0159 ^{NS} | 0.0069 ^{NS} |
| Error | 24 | 0.0019 | 0.0072 | 0.0034 |

*, ** significant at the 5% and 1% levels of probability, NS = Nonsignificant

Table 7. Value of the combining power of sorghum diallel crosses.

| Female/Male | Lamongan 1 | Tulungagung 2 | 'Jombang' |
|---------------------------------|------------|---------------|-----------|
| Seed weight | | | |
| Lamongan 1 | -0.0003 | -0.0053 | -0.0042 |
| Tulungagung 2 | -0.0045 | 0.0001 | -0.0027 |
| Jombang | -0.0003 | -0.0024 | 0.0002 |
| Seed vertical diameter | | | |
| Lamongan 1 | 0.2575 | 0.0610 | 0.0550 |
| Tulungagung 2 | 0.2042 | 0.2900 | 0.0700 |
| Jombang | 0.0782 | 0.1225 | 0.1825 |
| Seed horizontal diameter | | | |
| Lamongan 1 | -0.0028 | -0.0201 | -0.0326 |
| Tulungagung 2 | -0.0488 | 0.0164 | -0.0343 |
| Jombang | -0.0063 | -0.0137 | -0.0136 |

GCA effects: diagonal values, SCA effects: Values above the diagonal, Rec: Values below the diagonal

Table 8. Specific combining ability effects for seed weight and seed vertical and horizontal diameters in six F₁ diallel hybrids.

| F ₁ populations | Seed weight | Seed | |
|----------------------------|-------------|-------------------|---------------------|
| | | Vertical diameter | Horizontal diameter |
| Lamongan-1 × Tulungagung-2 | -0.0053 | -0.0458 | -0.0201 |
| Lamongan-1 × Jombang | -0.0042 | -0.0540 | -0.0326 |
| Tulungagung-2 × Lamongan-1 | -0.0045 | -0.0716 | -0.0488 |
| Tulungagung-2 × Jombang | -0.0027 | -0.0451 | -0.0343 |
| Jombang × Lamongan-1 | -0.0003 | -0.0116 | -0.0063 |
| Jombang × Tulungagung-2 | -0.0024 | -0.0262 | -0.0137 |

Cluster analysis

Cluster analysis for quantitative and qualitative characters was carried out by using Mini Tab software version-17. The results were presented in a dendrogram (Figure 1). In the cluster analysis, the sorghum parental genotypes were designated as G2 ('Lamongan-1'), G8 ('Tulungagung-2'), and G-9 ('Jombang'). Cluster analysis divided all of the parental genotypes and their F₁ hybrids into two large groups with a similarity of 13.13%. The first group consisted of five F₁ hybrids (G2 × G8, G8 × G2, G8 × G9, G9 × G2, and G9 × G8) and the three parental genotypes 'Lamongan-1', 'Tulungagung-2', and 'Jombang'. The second group comprised the F₁ hybrid G2 × G9. The first group consisted of two subgroups, which had a similarity of 42.28%. The first subgroup contained only one F₁ hybrid (G2 × G8), whereas the second subgroup consisted of four F₁ hybrids (G8 × G2, G8 × G9, G9 × G2, and G9 × G8) and the three parental sorghum cultivars 'Lamongan-1', 'Tulungagung-2', and 'Jombang'.

DISCUSSION

Seed setting is the result of seed formation at harvest after hybridization. The percentage of the success rate of crosses is measured as the number of seeds formed (seed set) divided by the number of flowers crossed multiplied by 100%. A cross is said to be successful if it is marked by the formation of ovules and unsuccessful if ovules are not formed as indicated by the drying of the husks. Syukur et al. (2010) opined that if the prospective fruit begins to enlarge and does not fall out, then fertilization has occurred. On the other hand, if the prospective fruit does not enlarge or fall out, then fertilization has failed.

The highest percentage of seed formation was observed in the F₁ hybrid 'Tulungagung-2' × 'Lamongan-1' (37.50%). The seed formation rates of the five other F₁ hybrids 'Lamongan-1' × 'Tulungagung-2', 'Lamongan-1' × 'Jombang', 'Tulungagung-2' × 'Jombang', 'Jombang' × 'Lamongan-1', and 'Jombang' × 'Tulungagung-2' were 4.38%, 3.57%, 4.89%, 7.26%, and 18.97%,




Figure 1. Dendrogram of the quantitative and qualitative traits of sorghum parental genotypes and their F₁ hybrids.

respectively. Many factors, including the condition of the pollen used, environmental factors, the receptive time of the female and male anthesis, and the level of the compatibility of the parental genotype used influence the success of crosses in sorghum. Pollen is highly susceptible to drought stress and thus quickly loses viability in accordance with the opinion of Lansac *et al.* (1994) and Andriani and Isnaini (2010), who stated that pollen cannot recover their viability even after rehydration under drought stress conditions and can be viable only for 3–6 h after blooming. Environmental factors, such as rainfall, sunlight, temperature, and humidity, can also affect the occurrence of fertilization. Syukur *et al.* (2010) stated that rainfall and high temperature cause the low success of hybridization in conventional crosses.

In sorghum, floral reception and anthesis is the period when flowers mature and become ready for pollination. In sorghum, female flowers are receptive for 3 days after blooming. The pistil (stigma) of sorghum flowers begins to enter the receptive period 2 days before the flowers bloom and is receptive for up to 8 to 16 days after the flowers bloom; however, it is optimally receptive for 3 days after the flowers bloom (Stephens, 1934; Pedersen *et al.*, 1998; Bello, 2008). During pollination, when the female flowers have entered the receptive period, which is marked by the opening of husks in the morning, the husks on the flowers open for 20 min and close

completely 2 h later. In sorghum, the time of the anthesis of male flowers occurs before 10 am. Field observations have shown that pollen could be obtained at 5 am. The past findings of Stephens (1934) showed that sorghum flowers generally bloom (anthesis) before 10 am. Hermaphrodite flowers bloom first, and single flowers bloom 2–4 days later.

Cross compatibility is the capability of the plants to form fruit (Syukur *et al.*, 2010). Plant crosses that produce fruit are compatible, whereas those that do not produce fruit are incompatible. The level of the compatibility (matching) of the parental genotypes used is very distant. Kinship can affect the success of a cross, wherein the farther the kinship between the two parental genotypes used in the cross, the smaller the chance of obtaining F₁ plants. This phenomenon is in accordance with the opinion of Syukur *et al.* (2010), who stated that the implementation of crossing in distant relatives is difficult because of natural obstacles, such as weak hybrid seeds that cannot survive and sterile F₁ plants. These barriers lead to the low success rates of distant relative crosses. The success of the cross (crossing ability) in some cross combinations is limited by the capability to form seeds (seed setting), which is dependent on the genomes of both parents. Differences in the parental genome cause inhibition during and after fertilization. Martin (1970) stated that a cross is classified as compatible if it can produce fruit at rates above 20%.

The maximum and at-par seed weight were shown by the F₁ hybrid 'Jombang' × 'Tulungagung-2' (0.028 g) and the parental genotype 'Tulungagung-2' (G8, 0.028 g), followed by 'Lamongan-1' (G2, 0.025 g) and 'Jombang' (G9, 0.023 g). The genetic influence of the two parents is suspected to play a role in the result of the cross. Kamil (1979) and Panjaitan *et al.* (2015) explained that plant genes affect the height and weight of seeds, seed shape, and seed size. The F₁ hybrid 'Lamongan-1' × 'Tulungagung-2' (0.008 g) had lower seed weight than the parental genotypes 'Lamongan-1' (G2 - 0.025 g) and 'Tulungagung-2' (G8 - 0.028 g) likely due to imperfect fertilization given that a successful distant relative cross has a low chance of forming a seed set in accordance with the opinion of Syukur *et al.* (2010) and Rao *et al.* (2016), who explained that the farther the kinship between the two parental genotypes used in a cross, the smaller the chance of obtaining normal F₁ plants.

The analysis of seed vertical and horizontal diameter showed nonsignificant differences among sorghum genotypes likely as a result of the influence of the genetic makeup of the genotypes in accordance with the opinion of Lakitan (1995) and Supriono *et al.* (2017), who stated that in certain plants, seed size is generally not influenced by the environment but is more controlled by genetic factors. Genotypes showed nonsignificant differences for variables, i.e., live, normal, and abnormal sprouts. However, the normal and live sprouts were more pronounced in the F₁ hybrid 'Tulungagung-2' × 'Lamongan-1' than in other crosses, due to which the said hybrid produced a number of seeds. The germination of the seeds of parental genotypes and their F₁ hybrids is influenced by internal factors (seed moisture content and seed physical structure) and external factors (environmental factors, such as seed growth). Seed weight and seed size also affect seed germination. Large seeds have extensive food reserves with large embryo sizes such that the growth rate of germination is optimal. Satwiko (2013) explained that large and small seeds exhibit differences in the plant growth process and that compared with large seeds, small seeds have less optimal growth due to lower food reserves and smaller embryo size.

The germination percentage of the parental genotypes and their F₁ hybrids ranged from 66.67% to 100%. However, the parental genotypes and the F₁ hybrids 'Lamongan-1' × 'Tulungagung-2', 'Tulungagung-2' × 'Jombang', and 'Lamongan-1' showed a low percentage of

germination likely due to their genetic makeup. Kartasapoetra (2003) and Juliantisa (2017), stated that the genetic factors that affect seed vigor and the basic patterns of germination and growth are genetically inherited and differ from one species to another. Gardner (1991) and Juliantisa (2017) explained that seeds have low germination and vigor due to weakness; temperature fluctuations due to rain and sunlight that cause the chemical composition of seeds to change; and damage due to predator attacks that damage the crop and consequently reduce the quality and quality of the seeds.

Among genotypes, the F₁ hybrid 'Lamongan-1' × 'Jombang' had a higher plant height than its parental genotypes, i.e., 'Lamongan-1' and 'Jombang' likely as a result of the influence of genetic and environmental factors in accordance with the opinions of Gardner (1991) and Bunphan (2015), who stated that genetic and environmental factors affect plant growth. In addition to plant height, the leaves are a component of plant growth. The F₁ hybrids 'Lamongan-1' × 'Jombang' and 'Jombang' × 'Tulungagung-2' showed more leaves than all other sorghum genotypes. Tall genotypes are presumed to have a high number of leaves in line with the research of Siswanto (2015), who found that plant height is positively and significantly correlated with the number of leaves.

In different crop plants, heterosis and heterobeltiosis are widely used by the breeders to compare F₁ hybrids with their parental genotypes. Utomo *et al.* (2018) reported that the occurrence of heterosis is caused by the expression of genes that are passed down by both parental genotypes to their offspring. On the basis of these speculations, heterotic effects can be used to determine whether the parental genotype can be used in crosses or not. In this study, the F₁ hybrid 'Lamongan-1' × 'Jombang' showed the highest positive heterotic effects compared with its mid- and better parents, followed by the F₁ hybrid 'Tulungagung-2' × 'Lamongan-1' with midparent heterosis for germination percentage. For seed weight, the F₁ hybrid 'Jombang' × 'Tulungagung-2' presented the highest positive heterotic effects as compared with its mid-parent. For the vertical and horizontal dimensions of the seeds, the F₁ hybrid 'Tulungagung-2' × 'Lamongan-1' showed a positive heterotic effect over its mid-parent.

The estimation of combining ability values is very useful for breeders in identifying the best crossbreeding parents to obtain

superior offspring. The difference in GCA and SCA values for each tested sorghum genotype could be attributed to the influence of several factors, i.e., the genotypes used in the cross, the observed characters, the model of the cross, and the combining ability method used (Rini *et al.*, 2017; Izzah *et al.*, 2019). The analysis of GCA, SCA, and reciprocal effects revealed that SCA effects were significantly different for the vertical and horizontal diameters of the seeds. The significant SCA variances indicated that the diversity of the cross combinations resulted in offspring that are better or worse than their parental genotypes (Budiyant *et al.*, 2016). The significantly different GCA and SCA for these three characters indicated the presence of additive and nonadditive gene action (Mishra *et al.*, 2008). Another report showed that the significance of GCA and SCA implies that additive and dominance effects contribute to the genetic control for the character (Sahid *et al.*, 2020).

In terms of GCA effects, the parental genotype 'Tulungagung-2' had a high positive GCA for seed weight and seed horizontal diameter, whereas the genotype 'Lamongan-1' had a high positive GCA for seed vertical diameter. These results indicated that the parental genotype 'Tulungagung-2' is the best general combiner for seed weight and seed horizontal diameter. Zare *et al.* (2011) reported that genotypes with positive GCA have good capability to combine with other genotypes for certain characters. In general, a cross between two parents with high GCA values produces hybrids with high SCA due to the interaction effect between positive × positive alleles (Hairmansis and Aswidinnoor, 2015). However, the present research showed that the parental genotype 'Tulungagung-2', despite being a good general combiner, did not produce F₁ hybrids with the highest positive SCA values, and all the hybrids revealed negative SCA effects for the studied traits.

Simamarta (2010) stated that the similarity distance is low if it ranges between 60% and 65%. Broadly speaking, the analysis of quantitative and qualitative character clusters revealed a similarity level of 13.13%. In the cluster analysis results presented in Figure 1, the F₁ hybrid 'Jombang' × 'Lamongan-1' had 89.49% similarity with the parental genotype 'Jombang' (G9), and the hybrid 'Jombang' × 'Tulungagung-2' had the similarity of 78.13% with the genotype 'Jombang' (G9) and the hybrid 'Jombang' ×

'Lamongan-1'. The F₁ hybrids 'Tulungagung-2' × 'Lamongan-1' and 'Tulungagung-2' × 'Jombang' had 65.28% similarity with the female parental genotype 'Tulungagung-2' (G8) likely due to the genetic makeup of the genotypes. Anderson and Joseph (2016) and Siswanto *et al.* (2015) opined that genetic makeup and environmental and growing conditions can affect the potential yield of the sorghum genotypes. Mangoendidjojo (2003) stated that the highest diversity could be obtained in crosses between distantly related genotypes.

CONCLUSIONS

2
The parental genotypes and their F₁ hybrids revealed significant differences in seed weight and seed diameter and nonsignificant differences in germination percentage and seed setting. The parental genotype 'Tulungagung-2' and its F₁ hybrids 'Jombang' × 'Tulungagung-2', 'Tulungagung-2' × 'Lamongan-1', and 'Jombang' × 'Tulungagung-2' exhibited the highest germination percentage, seed setting, seed weight, and seed diameter. 1
The F₁ hybrids 'Lamongan-1' × 'Jombang', 'Tulungagung-2' × 'Lamongan-1', 'Tulungagung-2' × 'Jombang', 'Jombang' × 'Lamongan-1', and 'Jombang' × 'Tulungagung-2' showed higher germination percentages of 100%, 88.00, 09%, 66.67%, 75.00%, and 83.33%, respectively, than the F₁ hybrid 'Lamongan-1' × 'Tulungagung-2', which had zero germination. The parental genotype 'Tulungagung-2' had the highest GCA effects and was identified as a good general paternal and maternal parent for the production of promising F₁ hybrids. The hybrid 'Jombang' × 'Lamongan-1' had the lowest negative SCA values for all 1
the characters and is thus recommended for improving grain yield. Cluster analysis divided all of the sorghum genotypes into two large groups with 13.13% similarity. The F₁ hybrid 'Jombang' × 'Lamongan-1' had the greatest similarity with the parental genotype 'Jombang' (G9, 89.49%).

5 ACKNOWLEDGEMENTS

The authors would like to thank the Directorate of Research and the Community Service University of Muhammadiyah Malang has funded this research through the center of study's flagship program in 2017.

REFERENCES

- Anderson FW, Joseph EK (2016). Yield Components in Hybrid Versus Inbred Sweet Sorghum. *Crop Science*.
- Andriani A, Isnaini (2013). Morphology and growth phase of sorghum. *Technological Innovation and Development*. IAARD, Press, Jakarta, Indonesia.
- Bello D (2008). Effect of pollination time on seed set in sorghum (*Sorghum bicolor* L. Moench) in Yola, Nigeria. *Res. J. Agron.* 2(3): 87-89.
- Boonlertnirun K, Srinives P, Sarithniran P, Jompuk C (2012). Genetic distance and heterotic pattern among single cross hybrids within waxy maize (*Zea mays* L.). *SABRAO J. Breed. Genet.* 44(2): 382-397.
- Budiyanti T, Sobir S, Wirnas D, Sunyoto S (2016). Combined power and gene action on fruit characters and yield of a half diallel population of five papaya genotypes (*Carica papaya* L.). *J. Hort.* <https://doi.org/10.21082/jhort.v25n4.2015.p278-293>.
- Bunphan D, Prasit J, Jirawat S, Joseph EK, Anderson FW (2015). Heterosis and combining ability of F₁ hybrid sweet sorghum in Thailand. *Crop Sci.* 55: 178-187.
- DG-TPH (2000). Director-General of TPH (DG-TPH). Directorate General of Food Crops and Horticulture. General Guidelines for Seed Quality Analysis. Directorate of Seed Development, Indonesia.
- Fehr WR (1987). Principle of cultivar development theory and technique. Vol. I, Macmillan Publication Company, New York, USA.
- Gardner FP (1991). Cultivation Plant Physiology. UI Press. Jakarta, Indonesia.
- Griffing B (1956). Concept of general and specific combining ability in relation to diallel crossing system. *Aust. Biol. Sci.* 9(4): 433 - 493.
- Hairmansis A, Aswidinnoor H (2015). Combined character filling gray rice varieties that carry neutral alleles at S-5 locus. *Zuriat*. <https://doi.org/10.24198/zuriat.v16i2.6774>.
- Hallauer AR, Miranda FJB (1988). Quantitative Genetics in Maize Breeding. Iowa State University Press, Ames, USA.
- ICRISAT (2004). Sorghum, a Crop of Substance. International Crops Research Institute for the Semi-Arid Tropics. Andhra Pradesh, India.
- Izzah NK, Wicaksono INA, Tresniawati C, Wardiana E (2019). Analysis of combination power and character correlation of the number of cocoa fruits in half diallel crosses. *J. Ind. Plants Refreshments*. <https://doi.org/10.21082/jtidp.v6n3.2019>. pp. 99-108.
- Jaikishan I, Rajendrakumar P, Hariprasanna K, Balakrishna D, Bhat BV, Tonapi VA (2019). Identification of differentially expressed transcripts at critical developmental stages in sorghum (*Sorghum bicolor* L. Moench) in relation to grain yield heterosis. *3-Biotechnol.* 9: 239. doi:10.1007/s13205-019-1777-6.
- Jimoh WLO, Abdullahi MS (2017). Proximate analysis of selected sorghum cultivars. *Bayero J. Pure Appl. Sci.* 10(1): 285-288.
- Juliantisa (2017). Seed vigor of four genotypes of sorghum (*Sorghum bicolor* L.) moench harvested at two different stages of maturity after twelve months of storage. Faculty of Agriculture, University of Lampung, Indonesia.
- Kamil J (1979). Seed Technology. Department of Agronomy. Faculty of Agriculture. Andalas University, Padang, Indonesia.
- Kartasapoetra AG (2003). Seed Processing Technology. Rineka Cipta. Jakarta, pp. 112.
- Krisnawati A, Adie MM (2011). Heterosis, heterobeltiosis and gene action of agronomic characters (*Glycine max* L. Merrill). *Biol. News* 10(6): 827-836.
- Lakitan B (1995). Hortikultura: Theory, cultivation and Post Harvest. PT. Raja Grafindo Persada. Jakarta. Hlm 219.
- Lansac AR, Sullivan CY, Johnson BE, Lee KW (1994). Viability and germination of the pollen of sorghum (*Sorghum bicolor* L. Moench). *Ann. Bot.* 74(1): 27-33.
- Maftuchah, Febriana L, Sulistyawati, Reswari HA, Septia ED (2021). Morphological diversity and heritability of nine local sorghum (*Sorghum bicolor* L.) genotypes in East Java. *Biodiversity* 22: 1310-1316.
- Maftuchah, Zainudin A, Reswari HA (2020). Seed production level, quality of crude Jatropha oil, and biodiesel Jatropha curcas as a biofuel resource. *J. Sci. Innov.* 20(3): 45-50.
- Mangoendidjojo W (2003). Plant Breeding. Kanisius. Yogyakarta.
- Martin JH (1970). History and Classification of Sorghum. In: Wall JS, Ross WM (Eds.). Sorghum Production and Utilization. The Avi Publishing Co. Inc. Westport CT, USA, pp. 702.
- Mishra TK, Panda RR, Bastia D, Bose LK (2008). Combining ability analysis of quantitative traits in rice bean (*Vigna umbrata* thunb.). *Emirates J. Food Agric.* <https://doi.org/10.9755/ejfa.v20i2.5190>.
- Panjaitan R, Zuhry E, Devino (2015). Characterization and genetic relationship of 15 genotypes batan's sweet sorghum (*Sorghum bicolor* L. Moench). *JOM Faperta* 2(1): 1-13.
- Pedersen JF, Kaeppler HF, Andrews DJ, Lee RD (1998). Chapter 14. Sorghum In: Banga SS, Banga SK (Eds.). Hybrid cultivar development. Springer-Verlag. India, pp. 432-354.
- Poehlman JM (1983). Breeding Field Crops, 2nd Ed. AU Publ. Co. Westport. CT, USA.
- Rao VT, P. Reddy PS, Reddy BVS (2016). Combining ability for grain mold resistance in sorghum (*Sorghum bicolor* L. Moench). *SABRAO J. Breed. Genet.* 48(3): 240-246.

- 9 Reddy PS, Reddy BVS, Rao PS (2014). Genotype by sowing date interaction effects on sugar yield components in sweet sorghum (*Sorghum bicolor* L. Moench). *SABRAO J. Breed Genet.* 46(2): 241-255.
- 13 Ringo J, Onkware A, Mgonja M, Deshpande S, Rathore A, Mneney E, Gudu S (2015). Heterosis for yield and its components in sorghum (*Sorghum bicolor* L. Moench) hybrids in drylands and sub-humid environments of East Africa. *Aust. J. Crop Sci.* 9(1): 9-13.
- Rini EP, Was D, Trikoesoemaningtyas, Sopandie D (2017). Genetic analysis on agronomic and quality traits of sorghum hybrids in Indonesia. *SABRAO J. Breed. Genet.* 49(2): 192-200.
- 12 Sahid ZD, Syukur M, Maharjaya A (2020). Combining ability and heterotic effects of chili pepper (*Capsicum annum* L.) genotypes for yield components and capsaicin content. *SABRAO J. Breed. Genet.* 52(30): 390-401.
- Satwiko T, Lahay RR, Damanik BSJ (2013). Tanggapan pertumbuhan dan produksi beberapa varietas kedelai (*Glycine max* L.) terhadap perbandingan komposisi pupuk. *J. Online Agroekoteknol.* 1(4): doi:10.32734/jaet.v1i4.4446.
- Sembiring H, Subekti NA (2010). The World's Major Sorghum Producers. Cereal Crop Research Institute. South Sulawesi, Indonesia.
- Simamarta J (2010). Software Engineering. Yogyakarta: Andi Publisher.
- 33 Singh RK, Chaudhary BD (1985). Biometrical Methods in Quantitative Genetic Analysis. Kalyani Publishers, New Delhi, India.
- Siswanto T, Zuhri E, Nurbaiti (2015). Yield and fat content of several varieties of sorghum (*Sorghum bicolor* L. moench.) given several doses of phosphorus fertilizer. *JOM Faperta* Vol. 2 No.2. Universitas Riau, Indonesia.
- 14 Slepier DA, Poehlman JM (2006). *Breeding Field Crops*. Fifth Edition. Blackwell Publishing. pp. 297-315.
- 22 Steel RGD, Torrie JH, Dickey DA (1997). *Principles and Procedures of Statistics: A Biometrical Approach*, 3rd ed. McGraw Hill Book Co., Inc., New York, 42.
- Stephens JC (1934). Anthesis, pollination, and fertilization in sorghum. *J. Agric. Res.* 49(2): 123-136.
- Suarni (2012). Potential of Sorghum as a Functional Foodstuff. Cereal Crop Research Institute. South Sulawesi, Indonesia.
- 16 Subagio H, Aqil M (2013). *Development of Sorghum Production in Indonesia*. Cereal Crop Research Institute. South Sulawesi, Indonesia.
- 5 Sulistyawati, Roeswitawati D, Ibrahim JT, Maftuchah (2019a). Agronomic character of several local genotypes of sorghum (*Sorghum bicolor* L.) at East Java. *IOP Conf. Series: Mater. Sci. Eng.* 532. doi:10.1088/1757-899X/532/1/012023.
- Sulistyawati, Roeswitawati D, Ibrahim JT, Maftuchah (2019b). Genetic diversity of local sorghum (*Sorghum bicolor* L.) genotypes of East Java, Indonesia for agro-morphological and physiological traits. *Biodiversity* 20(9): 2503-2510.
- Supriono, Zuhry E, Nurbaiti (2017). Evaluation of yield and seed quality on some cultivars of sorghum (*Sorghum bicolor* L.) with different planting spaces. *JOM Faperta* 4(1): 1-15.
- 8 Susilowati SH, Saliem HP (2013). Sorghum Trade in World and Asian Markets and Prospects for Development in Indonesia. In: Sumarno, et al. (eds). *Sorghum: Technological Innovation and Development*. IAARD Press, Jakarta, Indonesia.
- Syukur M, Yuniarti R, Sujiprihari S (2010). Artificial Crossing Techniques. Department of Agronomy and Horticulture IPB, Bogor, Indonesia, pp. 87-98.
- Utomo FH, Kristanto BA, Kusmiyati E (2018). A cross of 4 soybean varieties (*Glycine max* L.) for the assembly of drought-resistant soybeans. *J. Agro. Complex* 2(1): 93-101.
- 38 Welsh JR (1981). *Fundamentals of Plant Genetics and Breeding*. John Wiley and sons. New York. Chichester. Brisbane. Toronto 693-703. <https://doi.org/10.5897/AJAR10.646>.
- 7 Xie N, Chen LN, Wong KM, Cui YZ, Yang HQ (2016). Seed set and natural regeneration of *Dendrocalamus membranaceus* Munro after mass and sporadic flowering in Yunnan, China. *PLoS ONE* 11(4): e0153845. <https://doi.org/10.1371/journal.pone.0153845>.
- 17 Zare M, Choukan R, Heravan EM, Bihamta MR, Ordookhani K (2011). Gene action of some agronomic traits in corn (*Zea mays* L.) using diallel cross-analysis. *Afr. J. Agric. Res.* 6(12): 1-11.

Combaining ability and heterosis in sorghum_Sabrao

ORIGINALITY REPORT

27%

SIMILARITY INDEX

25%

INTERNET SOURCES

14%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

| | | |
|---|---|----|
| 1 | sabraojournal.org Internet Source | 5% |
| 2 | discovery.researcher.life Internet Source | 2% |
| 3 | www.searca.org Internet Source | 2% |
| 4 | pr.hec.gov.pk Internet Source | 1% |
| 5 | biodiversitas.mipa.uns.ac.id Internet Source | 1% |
| 6 | eprints.umm.ac.id Internet Source | 1% |
| 7 | link.springer.com Internet Source | 1% |
| 8 | 1library.net Internet Source | 1% |
| 9 | academicjournals.org Internet Source | 1% |

| | | |
|----|---|------|
| 10 | pakbs.org Internet Source | 1 % |
| 11 | peerj.com Internet Source | 1 % |
| 12 | repository.uin-suska.ac.id Internet Source | 1 % |
| 13 | ugspace.ug.edu.gh:8080 Internet Source | 1 % |
| 14 | K. B. Choudhary, Vikas Khandelwal, Sheetal Raj Sharma. "Chapter 5 Sorghum Improvement: Male Sterility and Hybrid Breeding Approaches", Springer Science and Business Media LLC, 2022 Publication | <1 % |
| 15 | www.researchgate.net Internet Source | <1 % |
| 16 | www.smujo.id Internet Source | <1 % |
| 17 | etd.repository.ugm.ac.id Internet Source | <1 % |
| 18 | www.thejaps.org.pk Internet Source | <1 % |
| 19 | qdoc.tips Internet Source | <1 % |

20

Ketthaisong, Danupol, Bhalang Suriharn, Ratchada Tangwongchai, and Kamol Lertrat. "Combining ability analysis in complete diallel cross of waxy corn (*Zea mays* var. *ceratina*) for starch pasting viscosity characteristics", *Scientia Horticulturae*, 2014.

Publication

<1 %

21

www.acci.org.za

Internet Source

<1 %

22

www.wudpeckerresearchjournals.org

Internet Source

<1 %

23

Cyril Diatta, Thierry Klanvi Tovignan, Eyanawa Atchozou Akata, Beatrice Elohor Ifie et al. "Heterosis and combining ability for yield and resistance to grain mold in tannin-free photoperiod-insensitive sorghum [*Sorghum bicolor* (L) Moench] in Senegal", *Journal of Crop Science and Biotechnology*, 2021

Publication

<1 %

24

R. Darvishzadeh. "Genetic analysis of partial resistance to black stem (*Phoma macdonaldii*) in sunflower as measured by a seedling test", *Plant Breeding*, 6/2007

Publication

<1 %

25

www.bdigital.unal.edu.co

Internet Source

<1 %

26 T. M. El-Gazzar, M. M. Nada, A. H. Hussein, A. R. Dawood. "Evaluation of New Summer Squash Hybrids (*Cucurbita pepo* L.) Compare with some Commercial Cultivars", Journal of Plant Production, 2020
Publication <1 %

27 (4-5-14)
<http://203.158.6.22:8080/sutir/bitstream/123456789/34>
Internet Source <1 %

28 ec2-50-19-248-237.compute-1.amazonaws.com
Internet Source <1 %

29 jurnal.uns.ac.id
Internet Source <1 %

30 ejournal2.undip.ac.id
Internet Source <1 %

31 D S Hanafiah, Haryati, G M Damanik, M S Limbong, Joshua, F F Silaen. "Increased of soybean genetic variability through hand pollination", IOP Conference Series: Earth and Environmental Science, 2022
Publication <1 %

32 pdfs.semanticscholar.org
Internet Source <1 %

33 sabrao.org
Internet Source <1 %

| | | |
|----|--|------|
| 34 | www.cropj.com Internet Source | <1 % |
| 35 | Submitted to Mansoura University Student Paper | <1 % |
| 36 | nanopdf.com Internet Source | <1 % |
| 37 | "The Sorghum Genome", Springer Nature, 2016 Publication | <1 % |
| 38 | www.globalsciencebooks.info Internet Source | <1 % |
| 39 | www.semanticscholar.org Internet Source | <1 % |
| 40 | A. Elsayed, E. Elsaid, E. Elsherbiny. "THE PERFORMANCE OF LATE BLIGHT GENE Ph-3 IN TOMATO UNDER THE EFFECT OF LOCAL POPULATIONS FROM <i>Phytophthora infestans</i> ", Journal of Plant Production, 2016 Publication | <1 % |
| 41 | smujo.id Internet Source | <1 % |
| 42 | Kamala Venkateswaran, N. Sivaraj, S.R. Pandravada, M. Thirupathi Reddy, B. Sarath Babu. "Classification, Distribution and Biology", Elsevier BV, 2019 Publication | <1 % |

43

Vivek Thapliyal, Rajinder Kumar Dhall, Pooja Manchanda, Harleen Kaur. "Utilizing novel parthenocarpic gynoecious cucumber (*Cucumis sativus* L.) inbreds for exploiting their heterotic potential under poly-house conditions", Research Square Platform LLC, 2023

Publication

<1 %

44

Yan Guo, Pan Li, Nana Yuyama, Lubin Tan, Yongcai Fu, Zuofeng Zhu, Fengxia Liu, Chuanqing Sun, Hongwei Cai. "Identification of Quantitative Trait Locus for Seed Dormancy and Expression Analysis of Four Dormancy-Related Genes in Sorghum", Tropical Plant Biology, 2015

Publication

<1 %

45

Vidi Mercyana, Samanhudi, Puji Harsono. "The effect of biochar application and number of shoots on the growth of ratoon sorghum in vertisol soil", IOP Conference Series: Earth and Environmental Science, 2021

Publication

<1 %

46

repo.unand.ac.id

Internet Source

<1 %

47

M. Abed, E. Ibrahim, A. El-Shoura. "Development of Okra (*Abelmoschus esculentus* L. Moench) Hybrids Derived from

<1 %

Selected Inbreds under Drought Stress",
Journal of Plant Production, 2020

Publication

48

dr.lib.sjp.ac.lk

Internet Source

<1 %

49

Ali Shahzad, Hameed Gul, Muhammad Ahsan, Depeng Wang, Shah Fahad. "Comparative Genetic Evaluation of Maize Inbred Lines at Seedling and Maturity Stages Under Drought Stress", Journal of Plant Growth Regulation, 2022

Publication

<1 %

50

Cristián Jacob, Basilio Carrasco, Andrés R. Schwember. "Advances in breeding and biotechnology of legume crops", Plant Cell, Tissue and Organ Culture (PCTOC), 2016

Publication

<1 %

51

M. Sultan, M. Abdel-Moneam, S. Hammoud, A. Hefena. "ESTIMATES OF COMBINING ABILITY FOR SOME AGRO-MORPHOLOGICAL TRAITS ON SOME RICE GENOTYPES.", Journal of Plant Production, 2015

Publication

<1 %

52

Ojwang, P.P.O.. "Genetic analysis for resistance to bean fly (*Ophiomyia phaseoli*) and seed yield among common bean genotypes in a semi-arid environment", Field Crops Research, 20110131

<1 %

53

Sugeng Widodo, Joko Triastono, Dewi Sahara, Arlyna Budi Pustika et al. "Economic Value, Farmers Perception, and Strategic Development of Sorghum in Central Java and Yogyakarta, Indonesia", Agriculture, 2023

Publication

<1 %

54

academic.oup.com

Internet Source

<1 %

55

jebas.org

Internet Source

<1 %

56

joarps.org

Internet Source

<1 %

57

psa.net.pk

Internet Source

<1 %

58

researchspace.ukzn.ac.za

Internet Source

<1 %

59

www.regional.org.au

Internet Source

<1 %

60

"Estimation of Heterosis and Combining Ability Effects on Grain Yield and Some Agronomic Traits of Sorghum under Three NPK Fertilizers Levels", Egyptian Journal of Agronomy, 2016

Publication

<1 %

61 Adnan Rasheed, Muhammad Ilyas, Taj Naseeb Khan, Athar Mahmood et al. "Study of genetic variability, heritability, and genetic advance for yield-related traits in tomato (*Solanum lycopersicon* MILL.)", *Frontiers in Genetics*, 2023
Publication

62 Submitted to Higher Education Commission Pakistan
Student Paper

63 agry.um.ac.ir
Internet Source

64 chimie.utcluj.ro
Internet Source

65 core.ac.uk
Internet Source

66 dspace.nwu.ac.za
Internet Source

67 ejfa.me
Internet Source

68 www.repository.embuni.ac.ke
Internet Source

69 www.scilit.net
Internet Source

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off